Findling Surgical, P.C.

Fillip M. Findling D.O. Surgery of the Hand. Wrist. & Elbow

Long Island Office: 15 Munro Blvd. Valley Stream, New York 11581

Phone: (516) 791-5800 Fax: (516) 837-3999 Queens Office: 58-50 Catalpa Ave Ridgewood, New York 11385 Phone: (718) 418-HAND (4263)

Fax: (718) 821-1125

Hand Surgery Post Operative Instructions

DIET: Regular diet. Start light and progress as tolerated. No alcoholic beverages on the day of surgery.

ACTIVITY:

- 1. Keep hand/wrist above the level of your heart <u>at all times</u> for the next 7 days. This should be accomplished by using the **FOAM PILLOW**. A sling will not hold your hand/wrist above your heart and therefore is inadequate (it also may cause shoulder stiffness).
- 2. Attempt to move unsplinted fingers as much as possible (UNLESS TENDON OR NERVE REPAIR PERFORMED).
- 3. Avoid activities which may re-injure your hand or finger.
- 4. Move all joints of the extremity that are not immobilized (i.e. shoulder, elbow, fingers, and thumb unless instructed otherwise).

DRESSING/ WOUND:

- 1. Keep your dressing clean and **DRY**. Do not change your dressing.
- 2. You may shower tomorrow. Cover your bandage with a plastic bag and use tape or a rubber band so that it is water tight.
- 3. If plaster has been applied or a cast has been placed —do not make holes in it, remove it, or stick objects in it (i.e. coat hangers, pencils).
- 4. Use the **FOAM PILLOW** as much as possible to protect the operative site as well as keep it elevated

PAIN MEDICINE:

- 1. Take pain medicine on an "AS NEEDED" basis according to your doctor's instructions.
- 2. Your pain will decrease over the next few days-- allowing you to:
 - Decrease your pain medicine quantity until you stop.
 - Increase the time between doses until you stop.
- 3. You should not drink alcoholic beverages while on pain medication.
- 4. Take pain medicine with food to prevent nausea.
- 5. Constipation can occur. If no bowel movement occurs within 48 hours take a laxative of your choice (over the counter).

CONTACT PHYSICIAN FOR:

Slight pain, swelling and bluish discoloration are to be expected. If you have breathing difficulty or chest pain dial 911 immediately. However, if the following symptoms occur notify your physician:

- •Temperature above 101° F
- Uncontrolled nausea/vomiting
- Signs of wound infection (redness, swelling, pus-like drainage)
- Excessive swelling and tightness
- •Inability to urinate in 8 hours
- progressively increasing pain
- Excessive bleeding
- Increasing numbness

OFFICE APPOINTMENT:

Call to schedule an appointment wit	your doctor in	days	tor a o	dressing	ı change

Patient Name		Signature	Date
	Witness	Date	